

Search for 07/023, 483

Trying 3106016892...Open

```
Welcome to STN International! Enter x:x
LOGINID:ssspta1806jxt
PASSWORD:
TERMINAL (ENTER 1, 2, 3, OR ?):2
```

* * * * * * * * * * Welcome to STN International * * * * * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Sep 29 The Philippines Inventory of Chemicals and Chemical Substances (PICCS) has been added to CHEMLIST
NEWS 3 Oct 27 New Extraction Code PAX now available in Derwent Files
NEWS 4 Oct 27 SET ABBREVIATIONS and SET PLURALS extended in Derwent World Patents Index files
NEWS 5 Oct 27 Patent Assignee Code Dictionary now available in Derwent Patent Files
NEWS 6 Oct 27 Plasdoc Key Serials Dictionary and Echoing added to Derwent Subscriber Files WPIDS and WPIX
NEWS 7 Nov 29 Derwent announces further increase in updates for DWPII
NEWS 8 Dec 5 French Multi-Disciplinary Database PASCAL Now on STN
NEWS 9 Dec 5 Trademarks on STN - New DEMAS and EUMAS Files
NEWS 10 Dec 15 2001 STN Pricing
NEWS 11 Dec 17 Merged CEABA-VTB for chemical engineering and biotechnology
NEWS 12 Dec 17 Corrosion Abstracts on STN
NEWS 13 Dec 17 SYNTHLINE from Prous Science now available on STN
NEWS 14 Dec 17 The CA Lexicon available in the CAPLUS and CA files
NEWS 15 Jan 05 AIDSLINE is being removed from STN
NEWS 16 Feb 06 Engineering Information Encompass files have new names
NEWS 17 Feb 16 TOXLINE no longer being updated

NEWS EXPRESS FREE UPGRADE 5.0e FOR STN EXPRESS 5.0 WITH DISCOVER!
 (WINDOWS) NOW AVAILABLE

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS INTER General Internet Information

NEWS LOGIN Welcome Banner and News Items

NEWS PHONE Direct Dial and Telecommunication Network Access to STN

NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 12:44:10 ON 23 FEB 2001

=> file medline capplus biosis

COST IN U.S. DOLLARS

SINCE FILE TOTAL
ENTRY SESSION

FULL ESTIMATED COST

0.15

0.15

FILE 'MEDLINE' ENTERED AT 12:44:26 ON 23 FEB 2001

FILE 'CAPLUS' ENTERED AT 12:44:26 ON 23 FEB 2001
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 12:44:26 ON 23 FEB 2001
COPYRIGHT (C) 2001 BIOSIS(R)

=> s (quantitat? or quantif?)(10a)nucleic acid(10a)(PCR or polymerase chain reaction#)

L1 151 (QUANTITAT? OR QUANTIF?)(10A) NUCLEIC ACID(10A)(PCR OR
POLYMERIC
E CHAIN REACTION#)

=> s l1 and (nucleotide#(10a)binding)

L2 0 L1 AND (NUCLEOTIDE#(10A) BINDING)

=> s l1 and (dNTP(10a)bind?)

L3 0 L1 AND (DNTP(10A) BIND?)

=> s l1 and dNTP

L4 0 L1 AND DNTP

=> s l1 and nucleotide#

L5 20 L1 AND NUCLEOTIDE#

=> s l5 and binding species

L6 0 L5 AND BINDING SPECIES

=> s l5 and immobiliz?

L7 2 L5 AND IMMOBILIZ?

=> d 17 1-2 bib ab

L7 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2001 ACS

AN 1994:210031 CAPLUS

DN 120:210031

TI Amplification and detection process

IN Harris, Raymond John; Morris, Charles Phillip

PA University of South Australia, Australia; Adelaide Children's Hospital

SO PCT Int. Appl., 55 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

| | PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|----|------------|--|----------|-----------------|----------|
| PI | WO 9402634 | A1 | 19940203 | WO 1993-AU379 | 19930726 |
| | W: | AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP,
KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD,
SE, SK, UA, US, VN | | | |
| | RW: | AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,
BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG | | | |

EP 656068 A1 19950607 EP 1993-915557 19930726
EP 656068 B1 19991215
R: CH, DE, FR, GB, IT, LI, NL, SE
AU 698934 B2 19981112 AU 1993-45511 19930726
US 5849544 A 19981215 US 1995-374764 19950124

PRAI AU 1992-3705 19920724

WO 1993-AU379 19930726

AB A method for detecting a target nucleic acid sequence involves amplification and detection in the same vessel. The method carries out all steps in a single vessels, lowers the frequency of false-positives and

minimizes the spread of contaminants (no data). The target nucleic sequence is amplified in a vessel contg. an **immobilized** capture probe that may optionally become involved in the amplification reaction. A sample is incubated with the capture probe under conditions that allow the amplified target sequence to be bound by the capture probe, and the presence of bound target nucleic acid sequence is detd. A kit making use of the method is described. Any nucleic acid amplification method may be used in the amplification step. The method was demonstrated in optimization expts. to detect Mycoplasma fermentans using a 138 bp

section

of insertion sequence-like element as the target. The capture probe was **immobilized** on nitrated polycarbonate microtiter plate wells and the target sequence detected using asym. PCR with a biotinylated probe. Amplification products were quantified after capture using Europium-labeled avidin. The amplification was specific for M.

fermentans

with a lower limit of detection of 100-1000 organisms.

L7 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2001 ACS

AN 1993:185112 CAPLUS

DN 118:185112

TI Detection and quantification of nucleic acids and formation of labelled **immobilized** nucleic acids using a combination of DNA ligation and chain extension with DNA polymerase

IN Parton, Adrian

PA Scientific Generics Ltd., UK

SO PCT Int. Appl., 46 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

| | PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|----|------------|------|----------|-----------------|----------|
| PI | WO 9304199 | A2 | 19930304 | WO 1992-GB1526 | 19920819 |
| | WO 9304199 | A3 | 19930415 | | |

W: JP, US

RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE

PRAI GB 1991-17902 19910820

GB 1992-2962 19920212

AB A method for detection or quantitation of a nucleic acid in a sample that uses an **immobilized** probe-primer and a DNA polymerase and labeled **nucleotide**, or a DNA ligase and labeled oligonucleotide, is described. The sample is hybridized with the **immobilized** probe and the hybrid extended with DNA polymerase in the presence of labeled **nucleotides**. Alternatively, the probe-target complex is incubated with the labeled oligonucleotide which is complementary to part of the target sequence and the **immobilized** probe and labeled oligonucleotide are ligated together with DNA ligase. If the target sequence is present, a labeled, **immobilized** sequence will be produced. This method increases the specificity of the identification. If the target sequence is also amplified by PCR or ligase chain reaction before hybridization the sensitivity is also increased.

L7 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2001 ACS
AB . . . false-positives and minimizes the spread of contaminants (no data). The target nucleic sequence is amplified in a vessel contg. an immobilized capture probe that may optionally become involved in the amplification reaction. A sample is incubated with the capture probe under. . . to detect Mycoplasma fermentans using a 138 bp section of insertion sequence-like element as the target. The capture probe was immobilized on nitrated polycarbonate microtiter plate wells and the target sequence detected using asym. PCR with a biotinylated probe. Amplification products. . .

IT Genetic methods
(LCR (ligase chain reaction), single-tube method for nucleic acid detection and quantitation by hybridization and, immobilized capture probes in)

IT Genetic methods
(SNAAC (sequential nucleic acid amplification and capture), single-tube method for amplification and quantitation of nucleic acids, immobilized capture probes in)

IT Sex
(detn. of, single-tube method for nucleic acid detection and quantitation by hybridization, immobilized capture probes in, genotyping in)

IT Nucleic acid hybridization
(single-tube method for nucleic acid amplification and quantitation of nucleic acids by, immobilized capture probes in)

IT Plant breeding and selection
(single-tube method for nucleic acid detection and quantitation by amplification and hybridization in, immobilized capture probes in, genotyping in)

IT Polymerase chain reaction
(single-tube method for nucleic acid detection and quantitation by hybridization and, immobilized capture probes in)

IT Legal chemistry and medicine
(single-tube method for nucleic acid detection and quantitation by hybridization, immobilized capture probes in, for detn. of paternity or maternity)

IT Animal tissue
(typing of, single-tube method for nucleic acid detection and quantitation by hybridization, immobilized capture probes in, genotyping in)

IT Genetic methods
(NASBA (nucleic acid sequence-based amplification), Q.beta.-replicase-dependent, single-tube method for nucleic acid detection and quantitation by hybridization and, immobilized capture probes in)

IT Genetic methods
(NASBA (nucleic acid sequence-based amplification), single-tube method for nucleic acid detection and quantitation by hybridization and, immobilized capture probes in)

IT Taxonomy
(chemo-, nucleic acids in, single-tube method for nucleic acid detection and quantitation by hybridization, immobilized capture probes in, genotyping in)

IT Nucleotides, polymers
RL: BIOL (Biological study)
(oligo-, immobilized, blocked, as capture probes in SNAAC single-tube method for amplification and quantitation of nucleic acids)

=> d 17 2 kwic

L7 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2001 ACS

TI Detection and quantification of nucleic acids and formation of labelled **immobilized** nucleic acids using a combination of DNA ligation and chain extension with DNA polymerase

AB A method for detection or quantitation of a nucleic acid in a sample that uses an **immobilized** probe-primer and a DNA polymerase and labeled **nucleotide**, or a DNA ligase and labeled oligonucleotide, is described. The sample is hybridized with the **immobilized** probe and the hybrid extended with DNA polymerase in the presence of labeled **nucleotides**. Alternatively, the probe-target complex is incubated with the labeled oligonucleotide which is complementary to part of the target sequence and the **immobilized** probe and labeled oligonucleotide are ligated together with DNA ligase. If the target sequence is present, a labeled, **immobilized** sequence will be produced. This method increases the specificity of the identification. If the target sequence is also amplified by. . .

ST nucleic acid detn quantitation **immobilized** probe; PCR ligase chain reaction **immobilized** primer

IT **Polymerase chain reaction**
(in nucleic acid detection and quantitation using **immobilized** probes, increased sensitivity and specificity in relation to)

IT Genetic methods
(ligase chain reaction, in nucleic acid detection and quantitation using **immobilized** probes, increased sensitivity and specificity in relation to)

IT **Immobilization, biochemical**
(of nucleic acids, using **immobilized** hybridization probes, DNA polymerase and ligase in)

IT 9012-90-2, DNA polymerase 9015-85-4, DNA ligase

RL: USES (Uses)
(in nucleic acid detection and quantitation using **immobilized** probes, increased sensitivity and specificity in relation to)

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#) | [Search Form](#) | [Posting Counts](#) | [Show S Numbers](#) | [Edit S Numbers](#) | [Preferences](#)**Search Results -**

| Term | Documents |
|--|-----------|
| NUCLEOTIDE\$1 | 0 |
| NUCLEOTIDE.DWPI,EPAB,JPAB,USPT. | 40843 |
| NUCLEOTIDEA.DWPI,EPAB,JPAB,USPT. | 1 |
| NUCLEOTIDED.DWPI,EPAB,JPAB,USPT. | 6 |
| NUCLEOTIDEE.DWPI,EPAB,JPAB,USPT. | 1 |
| NUCLEOTIDEL.DWPI,EPAB,JPAB,USPT. | 1 |
| NUCLEOTIDEN.DWPI,EPAB,JPAB,USPT. | 1 |
| NUCLEOTIDES.DWPI,EPAB,JPAB,USPT. | 27527 |
| NUCLEOTIDEW.DWPI,EPAB,JPAB,USPT. | 1 |
| NUCLEOTIDEX.DWPI,EPAB,JPAB,USPT. | 1 |
| (L10 AND NUCLEOTIDE\$1).USPT,JPAB,EPAB,DWPI. | 3 |

[There are more results than shown above. Click here to view the entire set.](#)

Database:

- US Patents Full-Text Database
- JPO Abstracts Database
- EPO Abstracts Database
- Derwent World Patents Index
- IBM Technical Disclosure Bulletins

116 and bind\$

[Refine Search:](#)[Clear](#)**Search History****Today's Date: 2/23/2001**

| <u>DB Name</u> | <u>Query</u> | <u>Hit Count</u> | <u>Set Name</u> |
|---------------------|--|------------------|-----------------|
| USPT,JPAB,EPAB,DWPI | 110 and nucleotide\$1 | 3 | <u>L16</u> |
| USPT,JPAB,EPAB,DWPI | 110 and nucleotide\$1 | 3 | <u>L15</u> |
| USPT,JPAB,EPAB,DWPI | 110 and nucleotide\$1 | 0 | <u>L14</u> |
| USPT,JPAB,EPAB,DWPI | 110 and solid support\$1 | 1 | <u>L13</u> |
| USPT,JPAB,EPAB,DWPI | 110 and bind\$ specie\$1 | 0 | <u>L12</u> |
| USPT,JPAB,EPAB,DWPI | 110 bind\$ specie\$1 | 0 | <u>L11</u> |
| USPT,JPAB,EPAB,DWPI | (quantif\$ or quantitat\$) near5 nucleotide\$1 near5
(PCR or polymerase chain reaction\$1) | 3 | <u>L10</u> |
| USPT,JPAB,EPAB,DWPI | (quantif\$ or quantitat\$) near5 (PCR or polymerase
chain reaction\$1) | 1433 | <u>L9</u> |
| USPT,JPAB,EPAB,DWPI | (quantif\$ or quantitat\$) near25 nucleotide\$1 near25
bind\$ near25(PCR or polymerase chain reaction\$1) | 0 | <u>L8</u> |
| USPT,JPAB,EPAB,DWPI | (quantif\$ or quantitat\$) and nucleotide\$1 and bind!
and (PCR or polymerase chain reaction\$1) | 6645 | <u>L7</u> |
| USPT,JPAB,EPAB,DWPI | (quantif\$ or quantitat\$) near20 nucleotide\$1 near20
bind! near20 (PCR or polymerase chain reaction\$1) | 0 | <u>L6</u> |
| USPT,JPAB,EPAB,DWPI | (quantif! or quantitat!) near20 nucleotide\$1 near20
bind! near20 (PCR or polymerase chain reaction\$1) | 0 | <u>L5</u> |
| USPT,JPAB,EPAB,DWPI | (quantif! or quantitat!) near10 nucleotide\$1 near10
bind! near10 (PCR or polymerase chain reaction\$1) | 0 | <u>L4</u> |
| USPT,JPAB,EPAB,DWPI | (quantif! or quantitat!) near5 nucleotide\$1 near5
bind! near5 (PCR or polymerase chain reaction\$1) | 0 | <u>L3</u> |
| USPT,JPAB,EPAB,DWPI | (quantiif! or quantitat!) near10 nucleotide\$1 near10
bind! near10 (PCR or polymerase chain reaction\$1) | 0 | <u>L2</u> |
| USPT,JPAB,EPAB,DWPI | (quantiif! or quantitat!) near5 nucleotide\$1 near5
bind! near5 (PCR or polymerase chain reaction\$1) | 0 | <u>L1</u> |